



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GROUP THEORY.

In view of the interest that is being taken in America in the Theories of Finite and Continuous Groups as well as in their various applications it is thought desirable to have a special department in THE AMERICAN MATHEMATICAL MONTHLY devoted to problems in this field. The receipt of letters from a number of our contributors suggesting this innovation leads us to hope it will be a successful one. THE EDITORS.

1. Proposed on p. 21 as Miscellaneous and solved on p. 98, as Group Theory No. 1.

2. Proposed by W. BURNSIDE, The Croft, Bromley Road, Catford, England.

Show that the group of the biquadratic equation $x^4 + 2ax^2 + b = 0$, in which a and b are rational numbers, while $a^2 - b$ is not the square of a rational number, is in general a dihedral group of order 8; but that (i) if b is the square of a rational number the group is a non-cyclical Abelian group of order 4; and (ii) if $b = a^2 \div (1 + n^2)$, where n is a rational number, the group is a cyclical group of order 4.

3. Proposed by L. E. DICKSON, Ph. D., The University of Chicago

Show that the equation $x^4 - ax^3 + bx^2 - ax + 1 = 0$, in which a and b are rational numbers, while neither $2 - b + (\frac{1}{2}a)^2$ nor $(1 + \frac{1}{2}b)^2 - a^2$ is the square of a rational number, is irreducible in the domain R of rational numbers; determine its group for this domain.

NOTES.

Editor Dickson has been granted leave of absence for six months from the University of Chicago to accept the appointment of Research Assistant to the Carnegie Institution. Dr. Dickson's investigation will relate to the application of Group Theory to certain problems in Geometry and Function Theory. F.

School Mathematics announces the following list of associate editors: Robert J. Aley, University of Indiana; Joseph V. Collins, State Normal School at Stevens Point, Wis.; Clarence E. Comstock, Bradley Polytechnic Institute; Ellery W. Davis, University of Nebraska; David Felmley, State Normal School, Normal, Ill.; J. E. Gould, University of Washington; Franklin Turner Jones, University School, Cleveland, O.; G. A. Miller, Stanford University; W. F. Moncrief, Winthrop College; E. H. Moore, University of Chicago; Frank Morley, Johns Hopkins University; Charles W. Newhall, Shattuck School, Faribault, Minn.; H. B. Newson, University of Kansas; John C. Packard, High School, Brookline, Mass.; Arthur Schultze, New York High School of Commerce; Monroe B. Snyder, Central High School, Philadelphia; David Eugene Smith, Teachers College, Columbia University.